



**Table 43.** Frequency of occurrence and duration of hypoxia at NERR sites, 1996-1998.


Region	N	%N	<4 h	4-8 h	8-12 h	12-16 h	16-20 h	20-24 h	>24 h
West Coast	307	40	0.60	0.26	0.11	0.02	0.01	0.00	0.00
Northeast	90	12	0.87	0.07	0.04	0.00	0.01	0.00	0.01
Mid-Atlantic	125	16	0.73	0.21	0.03	0.02	0.00	0.00	0.01
Southeast	78	10	0.83	0.14	0.01	0.00	0.00	0.00	0.01
Gulf/Carib.	165	22	0.69	0.16	0.06	0.02	0.03	0.01	0.03
	765	Mean	0.75	0.17	0.05	0.01	0.01	0.00	0.01




**Figure 198.** Predicted annual occurrence and duration of hypoxia at West Coast NERRs.




**Figure 199.** Predicted annual occurrence and duration of hypoxia at Northeast Coast and Great Lakes NERRs.



**Figure 200.** Predicted annual occurrence and duration of hypoxia at Mid-Atlantic NERRs.



**Figure 201.** Predicted annual occurrence and duration of hypoxia at Southeast Coast NERRs.



**Figure 202.** Predicted annual occurrence and duration of hypoxia at Gulf of Mexico and Caribbean NERRs.